


Abstract

5 Compounds of the formula

Too 10X



I

10  
15  
20  
25  
30  
35

ps wherein R is lower-alkyl, R<sup>1</sup> is halogen, R<sup>2</sup> is C<sub>1-12</sub>-alkyl, R<sup>3</sup> is hydroxy, lower-alkoxy, lower-alkyl-carbonyloxy, lower-alkoxy-lower-alkylcar-bonyloxy, lower-alkylaminocarbonyloxy, arylaminocar-bonyloxy or aryl-lower alkylaminocarbonyloxy, X is C<sub>1-18</sub>-alkylene which optionally can be interrupted by 1,4-phenylene or interrupted or lengthened by 1,4-cyclohexylene, A is di- or tri-substituted 2-imidazolyl attached via an ethylene group or a substituted or unsubstituted heterocycle selected from the group consisting of benzimidazolyl, benzimidazolonyl, imidazo[4,5-c]pyridinyl, imidazo[4,5-c]pyridinonyl, benzthiazolyl, benzodiazepine-2,5-dion-1-yl and pyrrolo[2,1-c]-[1,4]benzodiazepine-5,11-dion-10-yl and n is the number 0 or 1, in the form of racemates and optical antipodes, as well as N-oxides and pharmaceutically usable acid addition salts thereof. The compounds of formula I have a pronounced calcium-antagonistic and anti-arrhythmic activity and can accordingly be used as medicaments, especially for the control or prevention of angina pectoris, ischaemia, arrhythmias, high blood pressure and cardiac insufficiency.